

# Care Partner Information

## *Alzheimer's Disease and Related Dementia Caregiving Tips*

### Home Safety Issues Part I: Dementia & Falls

Falls are common among older people, and can be very serious. Falls may cause fractures or other injuries, and people who fall often become very afraid of falling again. Falls can result in a decrease in mobility, loss of independence, and death. People with dementia are at a high risk of falls. Although falls can't always be prevented, there are things to do to decrease falls.

First, it's important to follow the general tips for preventing falls. Then, it's helpful to know how the changes of dementia make falls more common, and what to do to specifically reduce the risk of falls in people with dementia. Also, remember that a person's abilities will change over time as the dementia worsens.

#### **General Tips for Preventing Falls and Injuries**

Most falls in older adults are due to a combination of things, such as medications and balance problems. Therefore, most of the time, it takes several changes in order to prevent falls. The table below lists the things to do to reduce the risk of falls and injuries in all older adults.

<b>Risk of Falls and What to Do</b>	
Exercise	Encourage regular exercise, including strength and balance training. Community fall programs are available through your local area agency on aging.
Safe Home Environment	Keep areas well lit. Use night-lights. Remove small rugs, long cords, and other tripping hazards from floors and stairs. Put grab bars in the bathroom by the toilet and shower. A nurse or occupational therapist can help with a home safety check.
Strong Bones	Discuss calcium and vitamin D treatment with the doctor. Also, ask the doctor about osteoporosis.
Foot Care	Foot pain, long nails, and numb or burning feet can cause falls. Wear shoes in the home.
Eye Care	Poor vision can cause falls. See an eye doctor at least each year. Wear glasses if needed.
Medications	Many medications can cause dizziness, weakness, and poor balance. Talk with the doctor.
Changing Positions	Many people get lightheaded when standing up. Change positions slowly. Give the person time to adjust.

This Care Partner Information page is part of a series on caregiving tips. They are written to help family and community caregivers, direct care workers and community health representatives care for someone with Alzheimer's disease or other types of dementia. Available in English and Spanish at [www.aging.arizona.edu](http://www.aging.arizona.edu)

## Care Partner Information

### Specific Tips for Preventing Falls in People with Dementia

It is important to learn how and why the changes of dementia make falls more common. Then you will be more aware, understand what's going on, and know how to help decrease falls.

<b>Reasons for Falls in People with Dementia</b>	
<b>Condition or Symptom</b>	<b>How it May Increase Risk of Falling &amp; What to Do to Help</b>
Physical Weakness	They may be frail, not able to routinely exercise, or weak from a past stroke. If able, walking regularly can help maintain strength.
Changes in Gait and Balance	They may have a shuffling walk with poor coordination. It's good if you can have someone walk with them.
Poor Judgment	They may try to walk down the steps alone, or outside on ice, or in the heat. It's good to have someone walk with them.
Trouble with Space and Vision	They may be unable to judge steps or uneven flooring. Reduce clutter and obstacles in the area. Lock or limit access to unsafe areas.
Medication Side Effects	They may be taking many medications due to cognitive or behavioral problems. Talk with the provider to reduce certain medications with side effects that can drop blood pressure, increase confusion, worsen balance, and increase fall risk.
Restlessness	Usually from aches and pains, hunger, thirst, or the need to use the bathroom. They may be unable to tell others, and get up quickly and urgently putting themselves at risk. Offer activity (moving around), food, and water regularly. If they have pain, work with the provider to treat it.
Boredom or Loneliness	They may be wandering around looking for something to do. Offer activities and arrange for visitors. Consider installing locks out of sight to reduce wandering outside.

### Useful Websites

Fall Prevention Center of Excellence: <http://stopfalls.org/>

**Written by: Mindy J. Fain, MD, University of Arizona Center on Aging**

Alzheimer's disease and Related Dementia ~ Care Partner Information

Edited by an interprofessional team from the University of Arizona Center on Aging,  
Alzheimer's Association - Desert Southwest Chapter and Community Caregivers