
County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 48 (11/26/2023-12/2/2023), Issue 9



Key messages

- Influenza activity in Riverside County is moderate. Influenza A is the predominant virus during this season.
- Public Health Advisory for Influenza and Other Common Transmissible Respiratory Illnesses is available <u>here</u>.
- Riverside County Respiratory Illness (COVID-19, influenza, and RSV) Dashboard is available <u>here</u>.
- Early decline in vaccinations has been observed.
- Annual influenza vaccination is recommended for all persons aged ≥ 6 months who do not have contraindications.

Influenza (flu) is a contagious respiratory illness caused by influenza viruses that infect the nose, throat, and lungs. Some people, such as older people, young children, and people with certain health conditions, are at higher risk for serious flu complications. Based on the Centers for Disease Control and Prevention (CDC) Weekly Influenza Surveillance Report, the California Department of Public Health (CDPH) Influenza, RSV and Other Respiratory Viruses Weekly Report (Figures 1-2) and local positivity rate data (Figure 14), the current influenza-like illness (ILI) activity level in Riverside County is moderate^{1, 2}. Riverside County collects influenza data through a variety of sources, including CDC's Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE), California Reportable Disease Information Exchange (CaIREDIE), California Integrated Vital Records System (Cal-IVRS), California Immunization Registry (CAIR), Riverside County Public Health Laboratory, Riverside University Health System-Medical Center (RUHS-MC) and sentinel providers. This report summarizes the current influenza surveillance data in the county.




Figure 1. National ILI Activity Level Map from CDC, Week 48

Figure 2. State ILI Activity Level Map from CDPH, Week 48



Page 1 of 11 **Epidemiology & Program Evaluation** 4065 County Circle Drive, Riverside, CA 92503 http://www.rivcoph.org



County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 48 (11/26/2023-12/2/2023), Issue 9

Emergency Department Syndromic Surveillance

Emergency department (ED) data are retrieved from ESSENCE. ILI records were captured based on chief complaints and discharge diagnosis. During week 48, ILI accounted for 4.5% (N=773) of all ED visits in Riverside County (Figures 3-4). Comparing to the previous week, ILI-related ED visits increased by 0.2%. ILI-related ED visits occurred across all age groups (Figure 5). However, after adjustment for age³, children aged 0-4 accounted for 65.4% of all ILI-related ED visits during this influenza season (Figure 6).




Figure 4. Percentage of ILI-related ED Visits by Week

16% 2023-2024 Season 14% 2022-2023 Season 12% 10% 8% 1 5 6% 4% 2% 0% 40 41 42 43 44 45 46 47 48 49 50 51 52 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 Dec Feb Oct Nov Jan Mar Apr May Jun Jul Aug Sep





Figure 5. Percentage of ILI-related ED Visits by Age Group and Week

Figure 6. Cumulative Age-adjusted Proportion of ILI-related ED Visits by Age Group, 2023-2024



Riverside University

Page 2 of 11 **Epidemiology & Program Evaluation** 4065 County Circle Drive, Riverside, CA 92503 http://www.rivcoph.org




County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 48 (11/26/2023-12/2/2023), Issue 9

Emergency Department Syndromic Surveillance

In order to better understand the potential geographic differences in ILI activity level, a heat map was produced based on the percentage of ILI-related ED visits over the total ED visits by patient's zip code (Figure 7). Areas with higher ILI levels could be considered for additional outreach regarding prevention and mitigation of respiratory illnesses.










Outpatient Service Syndromic Surveillance

RUHS-MC reports ILI-related outpatient visits weekly, including any face-to-face, phone and video visits. ILI records were retrieved based on visit diagnosis, which was not laboratory-confirmed. During week 48, the percentage of outpatient visits attributed to ILI was 3.7% (N=320) with a decrease of 0.5% than the previous week (Figures 8-9). Similarly, ILI-related outpatients visits occurred across all age groups (Figure 10). After adjustment for age³, children aged 0-4 accounted for 57.7% of all ILI-related outpatient visits during this influenza season (Figure 11).






Figure 10. Percentage of ILI-related Outpatient Visits by Age Group and Week





Figure 11. Cumulative Age-adjusted Proportion of ILI-related ED Visits by Age Group, 2023-2024





Page 4 of 11 **Epidemiology & Program Evaluation** 4065 County Circle Drive, Riverside, CA 92503 http://www.rivcoph.org



Severe Influenza Cases and Influenza-Associated Outbreaks

CDPH requires mandatory reporting of laboratory-confirmed influenza-associated fatal pediatric cases <18 years of age. In addition, Riverside County Public Health requires mandatory reporting of influenza ICU and associated deaths in laboratory-confirmed cases for ages 0-64 years. Severe influenza case data were retrieved from CaIREDIE. To date, no influenza-associated fatal cases or ICU cases have been reported during this season (Table 1).

	Fatal Cases 0-17 Years	Fatal Cases 18-64 Years	ICU Cases 0-64 Years		
Week 48	0	0	0		
Season To Date	0	0	0		

CDPH requires mandatory reporting of any respiratory disease outbreak, including influenza. Outbreak is defined as two or more cases of ILI (from separate households) in a setting within a 72-hour period with at least one case of laboratory-confirmed influenza. ILI is defined as fever (>37.8°C or 100°F) and either cough or sore throat in the absence of a known cause other than influenza To date, no influenza-associated outbreaks have been reported during this influenza season.





Electronic Laboratory Reporting Summary

Per Title 17, California Code of Regulations section 2505, laboratory results for influenza are required to be reported, including all positive and non-positive (negative, indeterminate, etc.) test results from both nucleic acid amplification tests (NAAT) and non-NAAT diagnostic tests (e.g., high throughput antigen tests) are reportable within one day from facilities certified under CLIA to perform nonwaived (moderate- or high-complexity) testing. However, non-positive test results are not consistently reported by laboratories. Hence, positivity rates cannot be calculated based on the data from Electronic Laboratory Reporting (ELR). During week 48, 475 positive influenza laboratory results were reported with an increase of 29.3% compared with the previous week (Figure 12).




Figure 12. Number of Positive Influenza Laboratory Results by Week





Clinical Laboratory Surveillance

For a better estimate of positivity rate, influenza laboratory data from RUHS-MC are analyzed, which include all positive and nonpositive test results. During week 48, 362 specimens were tested and 37 of them (10.2%) were positive for influenza virus (Figures 13-14). Please note, a specimen positive for both influenza A and influenza B will be counted twice. Influenza A was the dominant strain, accounting for 89.2% of positive specimens. Between weeks 40 and 48, the cumulative positivity rate for influenza was 5.0%, lower than the California average of 5.7% but higher than the national average of $3.7\%^{2,5}$ (Table 2).






Figure 13. Number of Positive Influenza Specimens by Week

Figure 14. Positivity Rate by Week





	Riverside County		Calif	ornia²	Nationwide ⁵	
	Week 48	Week 48 Data Cumulative Since Week 40		Week 48 Data Cumulative Since Week 40		Data Cumulative Since Week 40
Specimens Tested	362	2,289	23,170	107,951	93,944	690,802
Positive Specimens	37 (10.2%)	115 (5.0%)	2,028 (8.8%)	6,190 (5.7%)	6,415 (6.8%)	25,243 (3.7%)
Positive Specimens by Type						
Influenza A	33 (89.2%)	100 (87.0%)	1,899 (93.6%)	5,754 (93.0%)	5,152 (80.3%)	18,772 (74.4%)
Influenza B	4 (10.8%)	15 (13.0%)	129 (6.4%)	436 (7.0%)	1,263 (19.7%)	6,471 (25.6%)






Public Health Laboratory Surveillance

Public Health Laboratory Surveillance is important for influenza virus characterization and early detection of novel viruses. Riverside County Public Health Laboratory actively requests positive specimens from clinical laboratories in Riverside County and monitors the proportion of circulating viruses that belong to each influenza subtype or lineage. During week 48, 0 positive specimen had been subtyped in Riverside County Public Health Laboratory (Table 3 and Figure 15).

	Riverside County		Calif	ornia²	Nationwide ⁵		
	Week 48	Data Cumulative Since Week 40	Week 48	Week 48 Data Cumulative Since Week 40		Data Cumulative Since Week 40	
A positive specimens tested	0	0	39	324	437	3,843	
A (H1) pdm09	0	0	28	246	218	2,656	
A (H3)	0	0	4	41	76	451	
A, not subtyped	0	0	7	37	143	736	
B positive specimens tested	0	0	4	29	110	935	
B Victoria	0	0	4	17	84	764	
B Yamagata	0	0	0	0	0	0	
B, not lineage typed	0	0	0	12	26	171	

Table 3. Influenza Specimens by Subtype or Lineage, This Week and Season to Date








Page 8 of 11 **Epidemiology & Program Evaluation** 4065 County Circle Drive, Riverside, CA 92503 http://www.rivcoph.org



Deaths Registered with Either or Both of Pneumonia and Influenza

Pneumonia and influenza (P&I) are among the leading causes of death in the United States, accounting for over 1.2% of all deaths in 2021⁶. During week 48, 9.1% (N=15) of deaths were due, in part, to P&I in Riverside County (Figures 15-16). Between weeks 40 and 48, the cumulative proportion of P&I death was 7.4%. P&I deaths in the recent weeks may be undercounted because of the lengthy death certificate processing time. Newly identified P&I deaths will be added to the according week. Overall, people aged 65+ accounted for the majority of P&I deaths in Riverside County and very few P&I deaths happened among young people aged 0-24 (Table 4). Pneumonia, rather than influenza, contributed to the vast majority of P&I deaths, which were largely affected by other pneumonia causing diseases such as SARS-CoV-2.



	Age 0-4	Age 5-24	Age 25-49	Age 50-64	Age 65+	Total
P&I Death	1	1	7	24	187	220
& Percent	(0.5%)	(0.5%)	(3.2%)	(10.9%)	(85.0%)	






Influenza Vaccinations

Influenza causes millions of illnesses, hundreds of thousands of hospitalizations and tens of thousands of deaths in the United States, but less than half of Americans get an annual flu vaccine. Flu vaccine distribution generally begins in August and continues until all of the vaccines are distributed. The date of first dose was used for those who received two doses of flu vaccine. By the end of week 48, 18.6% of residents in Riverside County (N=455,501) had received a flu vaccine, 1.9% lower than the corresponding cumulative coverage rate during the 2022-2023 influenza season (Figures 18-19). Vaccination coverage was highest among people aged 65+ and lowest among people aged 5-24 (Table 5). Only 82.2% of the records had valid race and ethnicity information. Latinx were largely underrepresented than their counterparts (Table 6). Please note currently providers are not required to submit flu vaccine data to CAIR. Therefore, the vaccinated population may be underestimated.








Figure 19. Influenza Vaccination Coverage among Riverside County Residents, August 2023-Present

Table 5. Riverside County Residents Vaccinated for Influenza by Age Group, August 2023-Present

	Age 0-4	Age 5-24	Age 25-49	Age 50-64	Age 65+	Total
Influenza Vaccination	22,174	71,023	90,808	101,023	170,473	455,501
& Percent	(15.6%)	(10.5%)	(11.2%)	(23.4%)	(43.7%)	(18.6%)





Table 6. Influenza Vaccinations by Racial/Ethnic Group in Riverside County, August 2023-Present								
	American Indian or Alaska Native	Asian	Native Hawaiian or Other Pacific Islander	Black or African American	White	Latinx	Multiracial	
Influenza Vaccination & Proportion	1,688 (0.5%)	38,926 (10.4%)	2,118 (0.6%)	23,429 (6.3%)	165,181 (44.1%)	133,621 (35.7%)	9,453 (2.5%)	
Proportion Based on Standard Population ³	0.5%	6.9%	0.3%	6.3%	32.2%	51.4%	2.5%	
Representation	Equal	Over by 3.5%	Over by 0.3%	Equal	Over by 11.9%	Under by 15.7%	Equal	

Influenza Vaccinations

Public Health Recommendations

The best way to reduce risk from seasonal flu infection and its potentially serious complications is to get vaccinated annually. Flu vaccines are designed to protect against the four viruses that will be most common this influenza season and are recommended for individuals 6 months and older. Flu vaccines for the 2023-2024 influenza season are available now. COVID-19 preventive measures, such as social distancing, hand washing, and mask wearing, can also effectively prevent influenza.

References

- 1. Weekly US Map: Influenza Summary Update: https://www.cdc.gov/flu/weekly/usmap.htm
- 2. Influenza, RSV and Other Respiratory Viruses Weekly Report: https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Immunization/Influenza.aspx
- 3. Riverside population estimates were retrieved from California Department of Finance: <u>https://dof.ca.gov/forecasting/demographics/projections/</u>
- 4. National and regional baseline information for outpatient illness surveillance: https://www.cdc.gov/flu/weekly/overview.htm
- 5. U.S. Virologic Surveillance: <u>https://www.cdc.gov/flu/weekly/index.htm#ClinicalLaboratories</u>
- 6. Mortality in the United States, 2021: <u>https://www.cdc.gov/nchs/data/databriefs/db456.pdf</u>



