







County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Indicators at-a-glance



Key messages

- Influenza activity in Riverside County is Moderate. Influenza A is the predominant virus during this season.
- Public Health Advisory for Influenza and Other Common Transmissible Respiratory Illnesses is available [here](#).
- Riverside County Respiratory Illness (COVID-19, influenza, and RSV) Dashboard is available [here](#).
- Lower vaccination coverage has been observed.
- Annual influenza vaccination is recommended for all persons aged ≥6 months who do not have contraindications.

Influenza (flu) is a contagious respiratory illness caused by influenza viruses that infect the nose, throat, and lungs. Some people, such as older people, young children, and people with certain health conditions, are at higher risk for serious flu complications. Based on the Centers for Disease Control and Prevention (CDC) Weekly Influenza Surveillance Report, the California Department of Public Health (CDPH) Influenza, RSV and Other Respiratory Viruses Weekly Report (Figures 1-2) and local positivity rate data (Figure 14), the current influenza-like illness (ILI) activity level in Riverside County is Moderate^{1,2}. Riverside County collects influenza data through a variety of sources, including CDC's Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE), California Reportable Disease Information Exchange (CalREDIE), California Integrated Vital Records System (Cal-IVRS), California Immunization Registry (CAIR), Riverside County Public Health Laboratory, Riverside University Health System-Medical Center (RUHS-MC) and sentinel providers. This report summarizes the current influenza surveillance data in the county.

Figure 1. National ILI Activity Level Map from CDC, Week 52





Figure 2. State ILI Activity Level Map from CDPH, Week 52



County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Emergency Department Syndromic Surveillance

Emergency department (ED) data are retrieved from ESSENCE. ILI records were captured based on chief complaints and discharge diagnosis. During week 52, ILI accounted for 7.8% (N=1,588) of all ED visits in Riverside County (Figures 3-4). Comparing to the previous week, ILI-related ED visits decreased by 1.2%. ILI-related ED visits occurred across all age groups (Figure 5). However, after adjustment for age³, children aged 0-4 accounted for 60.8% of all ILI-related ED visits during this influenza season (Figure 6).

Figure 3. Number of ILI-related ED Visits by Week




Figure 4. Percentage of ILI-related ED Visits by Week




Figure 5. Percentage of ILI-related ED Visits by Age Group and Week





Figure 6. Cumulative Age-adjusted Proportion of ILI-related ED Visits by Age Group, 2023-2024




County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Emergency Department Syndromic Surveillance

In order to better understand the potential geographic differences in ILI activity level, a heat map was produced based on the percentage of ILI-related ED visits over the total ED visits by patient's zip code (Figure 7). Areas with higher ILI levels could be considered for additional outreach regarding prevention and mitigation of respiratory illnesses.

Figure 7. Percentage of ILI-related ED Visits by Patient's Zip Code, Week 52



County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Outpatient Service Syndromic Surveillance

RUHS-MC reports ILI-related outpatient visits weekly, including any face-to-face, phone and video visits. ILI records were retrieved based on visit diagnosis, which was not laboratory-confirmed. During week 52, the percentage of outpatient visits attributed to ILI was 5.1% (N=478), with a decrease of 1.2% than the previous week (Figures 8-9). Similarly, ILI-related outpatient visits occurred across all age groups (Figure 10). After adjustment for age³, children aged 0-4 accounted for 55.0% of all ILI-related outpatient visits during this influenza season (Figure 11).

Figure 8. Number of ILI-related Outpatient Visits by Week




Figure 9. Percentage of ILI-related Outpatient Visits by Week




Figure 10. Percentage of ILI-related Outpatient Visits by Age Group and Week





Figure 11. Cumulative Age-adjusted Proportion of ILI-related ED Visits by Age Group, 2023-2024



County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Severe Influenza Cases and Influenza-Associated Outbreaks

CDPH requires mandatory reporting of laboratory-confirmed influenza-associated fatal pediatric cases <18 years of age. In addition, Riverside County Public Health requires mandatory reporting of influenza ICU and associated deaths in laboratory-confirmed cases for ages 0-64 years. Severe influenza case data were retrieved from CalREDIE. Methods used to identify influenza-associated deaths or ICU hospitalizations differ from those used to identify influenza-coded deaths presented in the Mortality Surveillance. Influenza-associated cases require laboratory confirmation of influenza while influenza-coded deaths on death certificates do not necessarily consider laboratory testing. Therefore, influenza-associated deaths reported here might not include the same individuals in influenza-coded deaths.

To date, one influenza-associated fatal case (0-64 years old) and two influenza-associated ICU hospitalizations (0-64 years old) have been reported during this season (Table 1).

CDPH requires mandatory reporting of any respiratory disease outbreak, including influenza. Outbreak is defined as two or more cases of ILI (from separate households) in a setting within a 72-hour period with at least one case of laboratory-confirmed influenza. ILI is defined as fever (>37.8°C or 100°F) and either cough or sore throat in the absence of a known cause other than influenza.

To date, one influenza-associated outbreak has been reported during this influenza season (Table 1).

Table 1. Severe Influenza Cases, This Week and Season to Date

	Fatal Cases 0-64 Years	ICU Cases 0-64 Years	Outbreaks
Week 52	0	1	0
Season To Date	1	2	1


County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Electronic Laboratory Reporting Summary

Per Title 17, California Code of Regulations section 2505, laboratory results for influenza are required to be reported, including all positive and non-positive (negative, indeterminate, etc.) test results from both nucleic acid amplification tests (NAAT) and non-NAAT diagnostic tests (e.g., high throughput antigen tests) are reportable within one day from facilities certified under CLIA to perform non-waived (moderate- or high-complexity) testing. However, non-positive test results are not consistently reported by laboratories. Hence, positivity rates cannot be calculated based on the data from Electronic Laboratory Reporting (ELR). During week 52, 1,160 positive influenza laboratory results were reported with a decrease of 27.8% compared with the previous week (Figure 12).

Figure 12. Number of Positive Influenza Laboratory Results by Week



County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Clinical Laboratory Surveillance

For a better estimate of positivity rate, influenza laboratory data from RUHS-MC are analyzed, which include all positive and non-positive test results. During week 52, 472 specimens were tested and 59 of them (12.5%) were positive for influenza virus (Figures 13-14). Please note, a specimen positive for both influenza A and influenza B will be counted twice. Influenza A was the dominant strain, accounting for 86.4% of positive specimens. Between weeks 40 and 52, the cumulative positivity rate for influenza was 10.7%, lower than the California average of 11.1% but higher than the national average of 7.5%^{2,5} (Table 2).

Figure 13. Number of Positive Influenza Specimens by Week




Figure 14. Positivity Rate by Week




Table 2. Specimens Tested for Influenza by Type, This Week and Season to Date

	Riverside County		California ²		Nationwide ⁵	
	Week 52	Data Cumulative Since Week 40	Week 52	Data Cumulative Since Week 40	Week 52	Data Cumulative Since Week 40
Specimens Tested	472	4,131	28,815	223,784	102,294	1,186,063
Positive Specimens	59 (12.5%)	440 (10.7%)	4,830 (16.8%)	24,872 (11.1%)	17,925 (17.5%)	88,881 (7.5%)
Positive Specimens by Type						
Influenza A	51 (86.4%)	394 (89.5%)	4,519 (93.6%)	23,346 (93.9%)	14,732 (82.2%)	70,485 (79.3%)
Influenza B	8 (13.6%)	46 (10.5%)	311 (6.4%)	1,526 (6.1%)	3,193 (17.8%)	18,396 (20.7%)

County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13


Public Health Laboratory Surveillance

Public Health Laboratory Surveillance is important for influenza virus characterization and early detection of novel viruses. Riverside County Public Health Laboratory actively requests positive specimens from clinical laboratories in Riverside County and monitors the proportion of circulating viruses that belong to each influenza subtype or lineage. During week 52, 6 positive specimens had been subtyped in Riverside County Public Health Laboratory (Table 3 and Figure 15).

Table 3. Influenza Specimens by Subtype or Lineage, This Week and Season to Date

	Riverside County		California ²		Nationwide ⁵	
	Week 52	Data Cumulative Since Week 40	Week 52	Data Cumulative Since Week 40	Week 52	Data Cumulative Since Week 40
A positive specimens	6	100	73	824	581	9,552
A (H1) pdm09	5	80	63	648	300	6,254
A (H3)	1	20	4	107	42	1,368
A, not subtyped	0	0	6	69	239	1,930
B positive specimens	0	0	7	65	127	2,087
B Victoria	0	0	6	43	43	1,692
B Yamagata	0	0	0	0	0	0
B, not lineage typed	0	0	1	22	27	395

Figure 15. Percentage of Influenza Strains by Subtype/Lineage, This Week and Season to Date



County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Mortality Surveillance

Pneumonia and influenza are among the leading causes of death in the United States, accounting for over 1.2% of all deaths in 2021⁶. Riverside County analyzes death certificate data from California Integrated Vital Records System (Cal-IVRS) for all pneumonia and/or influenza deaths occurring in the Riverside County. Deaths in the recent weeks may be undercounted because of the lengthy death certificate processing time. Newly identified deaths will be added to the according week. Pneumonia and/or influenza-coded (P&I) deaths are defined as deaths who had pneumonia and/or influenza (text or coded) noted in any cause of death field on the death certificate, including immediate cause, underlying cause and other significant conditions. The International Classification of Diseases (ICD-10) codes used for P&I are J09-J18.

During week 52, 8.5% (N=9) of deaths were P&I deaths in Riverside County (Figures 16-17). Between weeks 40 and 52, the cumulative proportion of P&I deaths was 8.1%. Overall, people aged 65+ accounted for the majority of P&I deaths in Riverside County and very few P&I deaths happened among young people aged 0-24 (Table 4).

Figure 16. Number of Pneumonia and/or Influenza-Coded Deaths by Week




Figure 17. Percentage of Pneumonia and/or Influenza-Coded Deaths by Week




Table 4. Percentage of Pneumonia and/or Influenza-Coded Deaths by Age Group, Season to Date

	Age 0-4	Age 5-24	Age 25-49	Age 50-64	Age 65+	Total
P&I Death & Percent	1 (0.3%)	2 (0.5%)	17 (4.7%)	47 (12.9%)	298 (81.6%)	365

County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Mortality Surveillance

Pneumonia, rather than influenza, contributed to the vast majority of P&I deaths, which were largely affected by other pneumonia causing diseases such as COVID-19. P&I death no longer measures the impact of influenza in the same way it had prior to the COVID-19 pandemic, making it difficult to monitor the impact of influenza on mortality. Although monitoring influenza-only coded deaths will underestimate the full impact of influenza mortality, this measure allows for tracking trends in the impact of influenza on mortality and is not as influenced by COVID-19 as P&I deaths. Influenza-coded deaths are defined as deaths who had influenza (text or coded) noted in any cause of death field on the death certificate, including immediate cause, underlying cause and other significant conditions. The International Classification of Diseases (ICD-10) codes used for influenza are J09-J11. Please note, we are not able to verify influenza-associated death (severe influenza case) as influenza-coded death until their death certificate becomes available.

During week 52, 4 influenza-coded deaths were newly identified. To date, 11 influenza-coded deaths have been identified during this influenza season (Figure 18 & Table 5).

Figure 18. Number of Influenza-Coded Deaths by Week




Table 5. Number of Influenza-Coded Deaths by Age Group, Season to Date

	Age 0-4	Age 5-24	Age 25-49	Age 50-64	Age 65+	Total
Influenza-Coded Death	0	0	1	0	10	11

County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Influenza Vaccinations

Influenza causes millions of illnesses, hundreds of thousands of hospitalizations and tens of thousands of deaths in the United States, but less than half of Americans get an annual flu vaccine. Flu vaccine distribution generally begins in August and continues until all of the vaccines are distributed. The date of first dose was used for those who received two doses of flu vaccine. By the end of week 52, 20.5% of residents in Riverside County (N=502,215) had received a flu vaccine, 2.2% lower than the corresponding cumulative coverage rate during the 2022-2023 influenza season (Figures 19-20). Vaccination coverage was highest among people aged 65+ and lowest among people aged 5-24 (Table 6). Only 81.3% of the records had valid race and ethnicity information. Latinx were largely underrepresented than their counterparts (Table 7). Please note currently providers are not required to submit flu vaccine data to CAIR. Therefore, the vaccinated population may be underestimated.

Figure 19. Number of Riverside County Residents Vaccinated for Influenza, August 2023-Present




Figure 20. Influenza Vaccination Coverage among Riverside County Residents, August 2023-Present




Table 6. Riverside County Residents Vaccinated for Influenza by Age Group, August 2023-Present

	Age 0-4	Age 5-24	Age 25-49	Age 50-64	Age 65+	Total
Influenza Vaccination & Percent	26,042 (18.4%)	82,413 (12.2%)	101,258 (12.5%)	110,772 (25.6%)	181,730 (46.6%)	502,215 (20.5%)

County of Riverside Weekly Influenza Surveillance Report

2023 CDC Disease Week 52 (12/24/2023-12/30/2023), Issue 13

Influenza Vaccinations

Table 7. Influenza Vaccinations by Racial/Ethnic Group in Riverside County, August 2023-Present

	American Indian or Alaska Native	Asian	Native Hawaiian or Other Pacific Islander	Black or African American	White	Latinx	Multiracial
Influenza Vaccination & Proportion	1,909 (0.5%)	42,247 (10.3%)	2,352 (0.6%)	25,884 (6.3%)	176,492 (43.2%)	150,174 (36.8%)	9,381 (2.3%)
Proportion Based on Standard Population ³	0.5%	6.9%	0.3%	6.3%	32.2%	51.4%	2.5%
Representation	Equal	Over by 3.4%	Over by 0.3%	Equal	Over by 11.0%	Under by 14.6%	Under by 0.2%

Public Health Recommendations

The best way to reduce risk from seasonal flu infection and its potentially serious complications is to get vaccinated annually. Flu vaccines are designed to protect against the four viruses that will be most common this influenza season and are recommended for individuals 6 months and older. Flu vaccines for the 2023-2024 influenza season are available now. COVID-19 preventive measures, such as social distancing, hand washing, and mask wearing, can also effectively prevent influenza.

References

1. Weekly US Map: Influenza Summary Update: <https://www.cdc.gov/flu/weekly/usmap.htm>
2. Influenza, RSV and Other Respiratory Viruses Weekly Report: <https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Immunization/Influenza.aspx>
3. Riverside population estimates were retrieved from California Department of Finance: <https://dof.ca.gov/forecasting/demographics/projections/>
4. National and regional baseline information for outpatient illness surveillance: <https://www.cdc.gov/flu/weekly/overview.htm>
5. U.S. Virologic Surveillance: <https://www.cdc.gov/flu/weekly/index.htm#ClinicalLaboratories>
6. Mortality in the United States, 2021: <https://www.cdc.gov/nchs/data/databriefs/db456.pdf>